

8pt

5. How to store this medicine

- Keep all medicines out of reach and sight of children.
- Do not store above 25°C. Store in the original packaging.
- Do not use after expiry date shown on carton and sachet.
- Return any unused medicine to your pharmacist for safe disposal.

6. Further information**What is in this medicine****Active Ingredient**

Each 300mg granules in a sachet contain: Chamomilla recutita (Chamomilla) 6c

Also contains

Lactose monohydrate, Xylitol, Starch, Pregelatinised maize starch

What does the medicine look like

Teetha® Teething Granules are fine white to off-white granules in a sachet. There are 24 sachets in a box.

National rules authorisation holder and manufacturer is:

A Nelson & Co Limited,
5-9 Endeavour Way, Wimbledon,
London, SW19 8UH, UK
NR 01175/0387

This leaflet is available on request in formats suitable for the blind and partially sighted, please telephone: +44 (0)20 8780 4200.

This leaflet was revised: 12/2016

3001037/4826

8 pt

A Homeopathic Medicinal Product

Teetha® Teething Granules

Chamomilla 6c

Please read all of this leaflet carefully before giving this medicine, it contains important information.

Please keep this leaflet; you might need to read it again.

8pt all main body

1. What is this medicine for
Teetha® Teething Granules are a homeopathic medicinal product used within the homeopathic tradition for the symptomatic relief of teething pain and the symptoms associated with teething which are sore & tender gums, flushed cheeks and dribbling.

9pt all numbered headings

2. Before giving this medicine

- This medicine may not be suitable for some babies. Ask your doctor or pharmacist before giving it to the baby if you are not sure.
- Do not give this medicine to the baby if he/she is allergic to any of the ingredients, see section 6.
- This product contains lactose. If you have been told by your doctor that your baby has an intolerance to some sugars, consult your doctor before using this medicine.

P2

- If the baby is being given any other medicines including medicines prescribed by your doctor, or any medicines you have bought for them, talk to your doctor or pharmacist before giving them this medicine.

3. How to use this medicine**For oral use****For babies from 3 months old****Directions**

- Detach one sachet from strip of 4.

Babies 3-6 months

With the baby in an upright or sitting position.

Use a spoon.

Slowly give the baby half a sachet at a time into the front of the baby's mouth. Check the granules have fully dissolved before giving the remaining half.

Babies over 6 months

With the baby in an upright or sitting position.

Slowly pour the contents of the sachet into the front of the baby's mouth a little at a time. Alternatively use a spoon. Ensure the granules fully dissolve in the baby's mouth.

- Use one sachet every 2 hours for a maximum of 6 doses during any 24 hour period.

P3

- Consult a doctor or qualified healthcare practitioner if the symptoms worsen or persist for more than 7 days or if any side effects occur.
- **Do not** use more sachets than the label/leaflet tells you to.
- **If you forget to give this medicine to the baby**, continue with the usual dose at the usual time. It does not matter if you have missed a dose. Do not give a double dose to make up for a missed dose.
- **if you accidentally give the baby too much** consult your doctor or pharmacist as soon as possible.

4. Possible side effects

This medicine is not expected to cause any side effects. However, consult a doctor or qualified healthcare practitioner if the symptoms worsen, or persist for more than 7 days or if any side effects occur.

Reporting of side effects

If the baby gets any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the yellow card scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.